

Constant Domain Quantified Modal Logics Without Boolean Negation

Greg Restall

Greg.Restall@mq.edu.au

<http://www.mq.edu.au/~phildept/staff/grestall/>

Department of Philosophy, Macquarie University

Version of July 22, 1998

There is a natural way to add rules for first-order quantifiers to proof theories for propositional relevant logics, and there is a natural way to add evaluation conditions for quantifiers to the semantics for propositional relevant logics. Kit Fine has shown us that these two natural ways of modelling first order relevant logics do not match up [6].

Furthermore, Fine has given a semantics for the natural proof-theory for quantified relevant logics [5]. Unfortunately, the semantics is quite baroque. This has meant that the semantics has not produced any worthwhile results: The semantics has not been used to prove any interesting metatheoretical results about quantified relevant logics. Furthermore, the semantics has not been given anything like a plausible philosophical justification, or any new understanding of the unique features of quantified relevant logics.

In this paper I will begin to explore constant domain quantified relevant logics by examining a simpler case — I will look at constant domain *modal* logics without Boolean negation. That is, I will examine the proof theory and semantics for logics extending distributive lattice logic with extra *one place* operators. These are a simplification of relevant logics, which extend distributive lattice logic with the *two place* intensional connectives of implication and fusion. I end this paper by gesturing towards what needs to be done to extend the results of this paper to the more general setting of relevant logics.¹

1 Propositional Logic

I will motivate the *propositional* fragment of our logic semantically. We are interested in the relation of consequence defined on *frames*.

DEFINITION 1 A frame is a partially ordered set $\langle P, \sqsubseteq \rangle$ equipped with a number of binary relations of *accessibility*.

The *extensional* fragment of the language is interpreted in the usual way. We have two propositional constants \top and \perp , and two binary connectives \wedge and \vee .

¹I dedicate this paper to Richard Sylvan. He had a deep and abiding interest in relevant logics, including the problem of adding quantifiers in a semantically plausible fashion. His 1980 paper was an important addition to research in the area, but unfortunately, it contains errors which seem insurmountable [12]. It seems fitting that in an ancestor of this paper I thought I had solved the problems of constant domain quantified relevant logics by proving a stronger version of my Lemma 4, only to find a hole in the ‘proof’. I am also indebted to both Bernard Linsky and Ed Zalta [9] and Max Cresswell [8] who have, in conversation and in print, convinced me that constant domain semantics are philosophically interesting and defensible.

The “true at” relation \Vdash between points and propositions is defined recursively.² We have a case for atomic propositions which determines the behaviour of the relation \sqsubseteq , and the other cases determine the truth of complex propositions in terms of their constituents.

DEFINITION 2 An *evaluation* relation satisfies the following clauses.

- If $x \sqsubseteq x'$ and $x \Vdash p$ then $x' \Vdash p$.
- $x \Vdash A \wedge B$ iff $x \Vdash A$ and $x \Vdash B$.
- $x \Vdash A \vee B$ iff $x \Vdash A$ or $x \Vdash B$.
- $x \Vdash \top$.
- $x \not\Vdash \perp$.

We do not have Boolean negation for we are interested in ensuring that points can be *ordered*. Conjunction, disjunction, \top and \perp preserve the *hereditary* property of evaluations. If $x \Vdash A$ and $x \sqsubseteq x'$ then $x' \Vdash A$ too. Boolean negation on points (defined by setting $x \Vdash \neg A$ iff $x \not\Vdash A$) destroys this property.

This much is straightforward. The logic becomes more interesting when we add operators which exploit the nature of frames. We do this by using binary relations on frames. We allow for *four* different kinds of modal operators, \Box and \Diamond are familiar *positive* operators (called *A* and *I* operators) and \sim and \frown are less familiar *negative* operators (called *E* and *O*) respectively. They are evaluated according to the following conditions.³

DEFINITION 3 These are the evaluation conditions for *A*, *E*, *I* and *O* operators.

- $x \Vdash \Box A$ iff for all y where xRy , $y \Vdash A$.
- $x \Vdash \sim A$ iff for no y where xCy , $y \Vdash A$.
- $x \Vdash \Diamond A$ iff for some y where xSy , $y \Vdash A$.
- $x \Vdash \frown A$ iff for some y where xDy , $y \not\Vdash A$.

In the presence of Boolean negation it suffices to define *one* of these kinds of operators (say \Box) and to define each of the others in terms of it (as \Diamond is of the form $\neg\Box\neg$, \sim of the form $\Box\neg$ and \frown of the form $\neg\Box$.) In the absence of Boolean negation this strategy may not work, so we will take all four sorts of operators as primitive.

The hereditary result on formulae works only under special conditions on each accessibility relation. If $x \Vdash \Diamond A$ and $x \sqsubseteq x'$ then to ensure that $x' \Vdash \Diamond A$ we need a condition relating S and \sqsubseteq . The requisite conditions for each operator are listed below.

- If $x'Ry'$ and $x \sqsubseteq x'$ then there is some $y \sqsubseteq y'$ where xRy .

²The expression “ $x \Vdash A$ ” may be read as “ x supports A ”, “ x forces A ”, “ x makes A true” or “ A is true at x ” according to taste.

³If you are familiar with the theory of the syllogism, you will note why *A*, *E*, *I* and *O* for the families of one-place connectives. \Box is defined by a clause of the form *all A is B*, \Diamond by one of the form *some A is B*, \sim by one of the form *no A is B*, and \frown by one of the form *some A is not B*. These are the four syllogistic forms.

- If $x'Cy$ and $x \sqsubseteq x'$ then there is some $y' \supseteq y$ where xCy' .
- If xDy' and $x \sqsubseteq x'$ then there is some $y \sqsubseteq y'$ where $x'Dy$.
- If xSy and $x \sqsubseteq x'$ then there is some $y' \supseteq y$ where $x'Sy'$.

Each of the conditions on accessibility relations connects the accessibility relation with the behaviour of the inclusion relation on the frame. The inclusion relation can be read as the relation in *increasing informativeness*. If $x \sqsubseteq y$ then all of the information given by x is also given by y . Then, thinking of xCy as *x is compatible with y* motivates the condition on C . Similarly, reading xRy as *those things necessary in x are true in y* , and xSy as *y is possible relative to x* , motivates the conditions for R and S . (The interpretation of D is left an exercise for the interested. See *Consequences* Chapter 11 [11] for more details.)

DEFINITION 4 If M is a model, we say that $A \vdash_M B$ (A entails B on M) just when for each point x , if $x \Vdash A$ then $x \Vdash B$. Similarly, if F is a frame, $A \vdash_F B$ if and only if $A \vdash_M B$ for each model M on F . Finally, if \mathfrak{F} is a class of frames, $A \vdash_{\mathfrak{F}} B$ if and only if $A \vdash_F B$ for each frame $F \in \mathfrak{F}$.

The interpretation of the extensional connectives $\wedge, \vee, \top, \perp$ ensures that they are standard distributive lattice operators.

DEFINITION 5 A binary relation \vdash on propositions is a *distributive lattice relation* iff the following conditions hold:

- It is transitive.
- The two-place connective of conjunction is the *greatest lower bound* for \vdash . That is, $C \vdash A \wedge B$ if and only if $C \vdash A$ and $C \vdash B$.
- The two-place connective of disjunction is the *least upper bound* for \vdash . That is, $A \vee B \vdash C$ if and only if $A \vdash C$ and $B \vdash C$.
- Conjunction and disjunction are tied together with the distribution laws: $A \wedge (B \vee C) \vdash (A \wedge B) \vee (A \wedge C)$.
- There are special propositions \top and \perp such that $A \vdash \top$ and $\perp \vdash A$ for each A .

Each of the forms of consequence, \vdash_M , \vdash_F and $\vdash_{\mathfrak{F}}$ are distributive lattice consequence relations.

The other modal operators satisfy simple conditions, which we list in another definition.

DEFINITION 6 These are the inference conditions for each kind of operator.

- The A operator conditions are the inference from $A \vdash B$ to $\Box A \vdash \Box B$, and the entailments $\Box A \wedge \Box B \vdash \Box(A \wedge B)$ and $\top \vdash \Box \top$.
- The E operator conditions are the inference from $A \vdash B$ to $\sim B \vdash \sim A$, and the entailments $\sim A \wedge \sim B \vdash \sim(A \vee B)$ and $\top \vdash \sim \perp$.

- The I operator conditions are the inference from $A \vdash B$ to $\Diamond A \vdash \Diamond B$, and the entailments $\Diamond(A \vee B) \vdash \Diamond A \vee \Diamond B$ and $\Diamond \perp \vdash \perp$.
- The O operator conditions are the inference from $A \vdash B$ to $\neg B \vdash \neg A$, and the entailments $\neg(A \vee B) \vdash \neg A \vee \neg B$ and $\neg \top \vdash \perp$.

It is not difficult at all to verify that each of these conditions describes the behaviour of the modal operators and the consequence relations \vdash_M , \vdash_F and $\vdash_{\mathfrak{F}}$.

These operators can be defined in pairs in the style of Dunn's *Gaggle Theory* [1, 2, 4]. For example, A and I operators go together. If \Box and \Diamond satisfy the two-way rule

$$\Diamond A \vdash B \text{ if and only if } A \vdash \Box B$$

then \Diamond is an I operator and \Box is an A operator. Similarly, if \sim and \neg satisfy the two-way rule

$$A \vdash \sim B \text{ if and only if } B \vdash \neg A$$

then they are both E operators. Finally, if \frown and \smile both satisfy the rule

$$\frown A \vdash B \text{ if and only if } \smile B \vdash A$$

then they are both O operators.

These rules characterise the consequence relation $\vdash_{\mathfrak{F}}$ of the class of *all* frames. Soundness is trivial to verify. For completeness we can construct a *canonical model* out of prime theories. If you are interested in the logic of a smaller class of frames, those in which the accessibility relations satisfy interesting properties, you show that the canonical model of such a logic is one of those frames. If you are unlucky and the canonical frame doesn't satisfy the extra conditions, you will need to do more work by constructing a different frame.

This kind of semantic story has very many ancestors. My approach here is most indebted to the work done in the semantics for relevant logics due to Routley and Meyer [13], with further generalisations to Dunn [1, 3].

We can extend our relation \vdash to relate *sets* of propositions, by setting $\Sigma \vdash \Delta$ to be true just when some finite conjunction $\bigwedge_i A_i$ of members of Σ entails some disjunction $\bigvee_i B_i$ of members of Δ . We also use the convention that $\Sigma \vdash \emptyset$ iff $\Sigma \vdash \perp$, and $\vdash \Delta$ iff $\top \vdash \Delta$, and we sometimes write these as ' $\Sigma \vdash$ ' and ' $\vdash \Delta$ ' respectively. (The top and bottom elements \top and \perp are the empty conjunction and disjunction respectively.) By the soundness and completeness results and compactness theorem for these logics, this definition of set-based consequence is equivalent to the obvious frame condition: $\Sigma \vdash_M \Delta$ if and only if each point x forcing *each* element of Σ also forces *some* element of Δ . There is much more that we could do to examine the propositional logics. But for now, we will look at extending the logics to deal with quantifiers.

2 Models for Quantifiers

Our aim is to study the addition of constant domain quantifiers to this breed of model. To add quantifiers to the language, we enrich our language with *predicates, constants, variables* and the quantifiers \forall and \exists . We will define

formulae in such a way that we allow *formulae* with free variables, but we keep *sentences* to be formulae in which every occurring variable is bound.

For a constant domain model, we have not only a frame, but a non-empty *domain* of objects D . Each n -place predicate F is interpreted by a set $\|F\|_x$ of n -tuples of D -elements at each point x in the frame. This satisfies the hereditary condition that if $x \sqsubseteq y$ then $\|F\|_x \subseteq \|F\|_y$.

Terms (either constants or variables) are interpreted by domain elements, independently of the point in the model. (All terms are *rigid*.) Constants c are interpreted by objects $\|c\| \in D$. Variables v are interpreted by an *assignment* a which maps variables onto objects. We interpret v by $\|v\| = a(v)$, where a is an assignment. In either case, if t is a term, we use ' $\|t\|_a$ ' to name its interpretation, even in the case of constants, where the assignment is irrelevant to the interpretation.

Then we interpret formulas by relativising truth: not only to points in the model, but instead to pairs of points and assignments. Given an assignment a , and a point x we have

- $x, a \Vdash Ft_1 \cdots t_n$ iff $\langle \|t_1\|_a, \dots, \|t_n\|_a \rangle \in \|F\|_x$

Then to define the quantifiers, we have the standard conditions:

- $x, a \Vdash \forall v A$ iff for each $d \in D$, $x, a[v := d] \Vdash A$.
- $x, a \Vdash \exists v A$ iff for some $d \in D$, $x, a[v := d] \Vdash A$.

where $a[v := d]$ is the v -variant of the assignment a , which varies from a only by assigning the variable v the value d .

The connectives are interpreted in the usual way, as is the relation of consequence, relative to models, frames and classes of frames. Our job is to investigate these relations of consequence. The first thing to note is that the standard introduction and elimination rules for quantifiers hold.

INTRODUCTION AND ELIMINATION LAWS: For any formula A with v free we have

$$\forall v A(v) \vdash_M A(c) \text{ and } A(c) \vdash_M \exists v A(v)$$

for any model M , where $A(c)$ is found by replacing the free occurrences of v in $A(v)$ by the name c . These laws follow immediately from the evaluations of \forall and \exists . More interestingly, we have

$$\begin{aligned} &\text{If } A \vdash_F B(c) \text{ and } c \text{ does not occur in } A \text{ then } A \vdash_F \forall v B(v). \\ &\text{If } A(c) \vdash B \text{ and } c \text{ does not occur in } B \text{ then } \exists v A(v) \vdash B. \end{aligned}$$

for any frame F . If $A \vdash_F B(c)$ then any interpretation in which $x, a \Vdash A$, we must have $x, a \Vdash B(c)$. Now $x, a[v := d] \Vdash B$ if and only if $x, a \Vdash' B(c)$ for the variant interpretation \Vdash' which varies from \Vdash only by setting $\|c\| = d$. So, $x, a \Vdash \forall v B(v)$ if and only if $x, a \Vdash' B(c)$ for every such variant \Vdash' . However, \Vdash' and \Vdash do not disagree on the formula A , as c does not occur in A , so if $x, a \Vdash B(c)$ (which we have assumed) we also have $x, a \Vdash' B(c)$ too, so we have our result. Similar reasoning verifies the case of the existential quantifier.

These rules ensure that if v is not free in A , $\forall v A$ and $\exists v A$ are equivalent to A . For $\forall v A \vdash A$, and since v is not free in A , $A \vdash A(c)$ for some c not in A (since $A(c)$ is the very same formula as A !) so $A \vdash \forall v A$.

These rules are standard and work for almost any notion of logical consequence with quantifiers. The next results begin to exploit the power of the constant domains.

DISTRIBUTIVE LAWS: The quantifiers distribute over conjunction and disjunction in the following way.

$$\forall v(A \vee B) \vdash_M \forall vA \vee \exists vB \quad \exists vA \wedge \forall vB \vdash_M \exists v(A \wedge B).$$

These results hold in classical quantificational logic, but the first does not in quantified intuitionistic logic. However, they are straightforward applications of the rules. We will not tarry to verify them here.

These rules have simpler cases when B does not contain a free v . Since in that case B is equivalent to $\exists vB$ and $\forall vB$ we have $\forall v(A \vee B) \vdash \forall vA \vee B$, and $\exists vA \wedge B \vdash \exists v(A \wedge B)$.

Intuitionistic logic contains the second of these laws but not the first. Neither of these laws follows from distributive lattice logic together with the introduction and elimination laws.⁴

The most interesting results show the interaction between the quantifiers and the extra operators. These results are jointly called the *Barcan Laws*.

BARCAN LAWS: For any model M and any formula A , we have

$$\begin{aligned} \Box \forall vA \dashv\vdash_M \forall v\Box A & \quad \Diamond \exists vA \dashv\vdash_M \exists v\Diamond A \\ \sim \exists vA \dashv\vdash_M \forall v\sim A & \quad \neg \forall vA \dashv\vdash_M \exists v\neg A \end{aligned}$$

These results commute a quantifier over a one-place operator. They essentially rely on the fact that the domain D of quantification stays constant from point to point. They are straightforward to verify. For example, $x, a \Vdash \sim \exists vA$ if and only if for no y where xCy does $y, a \Vdash \exists vA$. That is, iff for no y where xCy does $y, a[v := d] \Vdash A$ for some $d \in D$, and this just means that for every $d \in D$, for no y does $y, a[v := d] \Vdash A$. This is equivalent to saying that for every $d \in D$, $x, a[v := d] \Vdash \sim A$, which is simply $x \Vdash \forall v\sim A$ as desired. If the domain D switched from point to point, we could not reason like this. The other Barcan laws are verified in a similar fashion.

These conditions: the introduction/elimination laws, the distributive laws, and the Barcan laws, jointly give us a good picture the behaviour of the universal and existential quantifiers in constant domain models and frames. It is our job to show that these laws *determine* the behaviour of the quantifiers on constant domain frames. To do this, however, we need some more tools. Somewhat surprisingly, we need two extra connectives, defined on frames, in order to prove that these rules characterise the logic of constant domain quantification on frames.

⁴For a model in which the introduction and elimination rules hold, use frames with *decreasing* domains (if $x \sqsubseteq x'$ then $D(x) \supseteq D(x')$). Set a universally quantified statement to be true at x just when every instance is true at x . Existentially quantified sentences are true at x just when some instance is true *at some ancestor of x* . These are the dual conditions to intuitionistic quantifiers, and they verify the distribution of universal quantifiers over disjunction, but not existential quantifiers over conjunction.

3 New Connectives

It is well known that you can conservatively add a connective \supset to the language of distributive lattice logic, satisfying the condition

$$A \wedge B \vdash C \text{ if and only if } A \vdash B \supset C$$

This is intuitionistic implication. The semantics is reasonably straightforward. We require that frames be extended with a partial order \sqsubseteq , and that atomic evaluations be preserved up the order. We set $x \Vdash A \supset B$ iff for each y where $x \sqsubseteq y$, if $y \Vdash A$ then $y \Vdash B$.

Note that with this addition, the unary operator \neg defined by setting $\neg A$ to $A \supset \perp$ is an E operator. $A \supset \top$ and $\perp \supset A$ are both equivalent to \top , and $\top \supset A$ is equivalent to A .

It's less well known that it's just as easy to extend things in the other direction. We can conservatively add a connective $-$ satisfying the condition

$$A \vdash B \vee C \text{ if and only if } A - B \vdash C$$

(You can read ' $A - B$ ' as ' A without B '.) The clause is the exact dual of that for \supset . For each x , $x \Vdash A - B$ iff there is some $y \sqsubseteq x$ where $y \Vdash A$ and $y \not\Vdash B$. This preserves the hereditary condition, and so, is a conservative extension of intuitionistic logic and therefore, of distributive lattice logic.

Note too that \frown , defined by setting $\frown A$ as $\top - A$ is an O operator. $\perp - A$ and $A - \top$ are both equivalent to \perp , while $A - \perp$ is equivalent to A .

You get conditions tying quantifiers to the conditional and to subtraction.

$$\forall v(A \supset B) \vdash_M \forall vA \supset \forall vB \quad \exists vB - \exists vA \vdash_M \exists v(B - A)$$

These can be verified semantically in a straightforward fashion. More interesting is the fact that they follow given the proof theory already at hand. We take the subtraction case as it is less familiar.

$$\begin{array}{c} B(c) - A(c) \vdash B(c) - A(c) \\ \hline B(c) - (B(c) - A(c)) \vdash A(c) \\ \hline B(c) - (B(c) - A(c)) \vdash \exists vA \\ \hline B(c) - \exists vA \vdash B(c) - A(c) \\ \hline B(c) \vdash \exists vA \vee \exists v(B - A) \\ \hline \exists vB \vdash \exists vA \vee \exists v(B - A) \\ \hline \exists vB - \exists vA \vdash \exists v(B - A) \end{array}$$

Furthermore, we have the "flipping" rules

$$\forall v(A \supset B) \vdash \exists vA \supset B \quad B - \forall vA \vdash \exists v(B - A)$$

where v is not free in B . These are verified similarly. Here is the subtraction case.

$$\begin{array}{c} B - A(c) \vdash B - A(c) \\ \hline B - A(c) \vdash \exists v(B - A) \\ \hline B - \exists v(B - A) \vdash A(c) \\ \hline B - \exists v(B - A) \vdash \forall vA \\ \hline B - \forall vA \vdash \exists v(B - A) \end{array}$$

Finally, and most interestingly, the introduction and elimination laws, together with implication and subtraction, is enough to verify the distribution laws.

$$\begin{array}{c}
A(c) \vee B(c) \vdash A(c) \vee B(c) \\
\hline
\forall v(A \vee B) \vdash A(c) \vee B(c) \\
\hline
\forall v(A \vee B) - A(c) \vdash B(c) \\
\hline
\forall v(A \vee B) - A(c) \vdash \exists vB \\
\hline
\forall v(A \vee B) - \exists vB \vdash A(c) \\
\hline
\forall v(A \vee B) - \exists vB \vdash \forall vA \\
\hline
\forall v(A \vee B) \vdash \forall vA \vee \exists vB
\end{array}$$

These connectives are definable on any constant domain frame using the structure already existing on that frame. Much more can be said about them (especially subtraction, which is much less studied than implication [10]). However, all of this is preliminary to our present project of adding constant domain quantifiers to our logic. It will become clear why having implication and subtraction at our disposal helps when we get to the completeness proof.

4 Completeness

For completeness we will construct a *canonical frame* and *model*. We will show that if $\Sigma \not\vdash \Delta$ (in the proof theory) then there is a point in the canonical model in which every element of Σ is true but every element of Δ is false. The completeness proof is not too different from that in either classical modal logic or that in propositional relevant logics. It is handy to define *pairs* of sets of formulae. The left member of a pair is the set of formulae we wish to make true and the right member of the pair is the set of formulae we wish to see fail.

DEFINITION 7 The pair $\langle \Sigma, \Delta \rangle$ is said to be a \vdash -*pair* (or simply, a *pair* in what follows) if $\Sigma \not\vdash \Delta$. A pair $\langle \Sigma, \Delta \rangle$ is *quantifier suited* iff the following two conditions hold:

- If $\Sigma \vdash \Delta \cup \{A(c)\}$ for each c then $\Sigma \vdash \Delta \cup \{\forall vA\}$ for each v .
- If $\Sigma \cup \{A(c)\} \vdash \Delta$ for each c then $\Sigma \cup \{\exists vA\} \vdash \Delta$ for each v .

A pair $\langle \Sigma, \Delta \rangle$ is said to be *full* in a language L iff $\Sigma \cup \Delta = L$.

Our canonical frame will be constructed from full quantifier-suited pairs from a particular language. The left set in such a pair is a good match for a point in a model. The next result is simple to prove.

LEMMA 1 *If $\langle \Sigma, \Delta \rangle$ is a full pair, then Σ is a prime theory. If $\langle \Sigma, \Delta \rangle$ is full and quantifier-suited, then $\exists vA \in \Sigma$ if and only if $A(c) \in \Sigma$ for some name c , and $\forall vA \in \Sigma$ if and only if $A(c) \in \Sigma$ for all names c .*

A prime theory satisfies each of the required distributive lattice conditions. If Σ is a prime theory, then $\top \in \Sigma$, $\perp \notin \Sigma$, $A \wedge B \in \Sigma$ if and only if $A, B \in \Sigma$, and $A \vee B \in \Sigma$ if and only if either A or B is in Σ . If the pair is also quantifier-suited, then the quantifiers are interpreted in the desired way too. $\forall vA \in \Sigma$ if and only if $A(c) \in \Sigma$ for each name c . $\exists vA \in \Sigma$ if and only if $A(c) \in \Sigma$ for some name c .

Now to *get* full quantifier-suited pairs. Firstly, we have the technique needed to get one in the first place.

LEMMA 2 (PAIR EXTENSION LEMMA 1) *If $\langle \Sigma, \Delta \rangle$ is a pair, then there is a full quantifier-suited pair $\langle \Sigma', \Delta' \rangle$ extending $\langle \Sigma, \Delta \rangle$, in a new language extending the original language by at most countably many new constants.*

PROOF This is achieved by enumerating the formulae in the language A_0, A_1, \dots and defining the pair $\langle \Sigma_0, \Delta_0 \rangle$ to be $\langle \Sigma, \Delta \rangle$, and $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle$ is defined as follows:

- If $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$ and A_n is not of the form $\exists vB$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n \cup \{A_n\}, \Delta_n \rangle$.
- If $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$ and A_n is of the form $\exists vB$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n \cup \{A_n, B(c)\}, \Delta_n \rangle$, where c is a new constant not appearing in Σ_n, A_n or Δ_n .
- If $\Sigma_n \cup \{A_n\} \vdash \Delta_n$ and A_n is not of the form $\forall vB$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n, \Delta_n \cup \{A_n\} \rangle$.
- If $\Sigma_n \cup \{A_n\} \vdash \Delta_n$ and A_n is of the form $\forall vB$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n, \Delta_n \cup \{A_n, B(c)\} \rangle$, where c is a new constant not appearing in Σ_n, A_n or Δ_n .

Then we set $\langle \Sigma', \Delta' \rangle$ to be $\langle \bigcup_n \Sigma_n, \bigcup_n \Delta_n \rangle$. This is a partition of the formulas by construction: every formula A_n is either in Σ' or Δ' .

We next show that it is a pair: We show that $\Sigma' \not\vdash \Delta'$. To show this, given the compactness of \vdash it suffices to show that $\Sigma_n \not\vdash \Delta_n$ for each n . We show this by induction on n . The result holds for $n = 0$ by hypothesis. For $n + 1$, suppose $\Sigma_{n+1} \vdash \Delta_{n+1}$ but that $\Sigma_n \not\vdash \Delta_n$. It follows that we cannot be in the first case of the construction, for that assures $\Sigma_{n+1} \not\vdash \Delta_{n+1}$ explicitly. Suppose A_n is of the form $\forall vB$, and that while $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$ we have $\Sigma_n \cup \{A_n, B(c)\} \vdash \Delta_n$ for a c new to Σ_n and Δ_n and A_n . Therefore there is a conjunction C of members of Σ_n , and a disjunction D from Δ_n such that $C \wedge A_n \wedge B(c) \vdash D$. It follows that $\exists v(C \wedge A_n \wedge B) \vdash D$ by existential introduction. By the existential distribution fact we have $C \wedge A_n \wedge \exists vB \vdash D$ (c is new to C, A_n and D) and hence $C \wedge A_n \vdash D$, contrary to our hypothesis that $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$. So, this case assures us that $\Sigma_{n+1} \not\vdash \Delta_{n+1}$.

Now suppose we are in the third case, and that both $\Sigma_n \cup \{A_n\} \vdash \Delta_n$ and that $\Sigma_n \vdash \Delta_n \cup \{A_n\}$. Then for some $C \in \bigwedge \Sigma_n$ and $D \in \bigvee \Delta_n$ we have $C \wedge A_n \vdash D$ and $C \vdash A_n \vee D$. Disjoining D to the first, we have $(C \wedge A_n) \vdash D$, and conjoining C to the second, we have $C \vdash C \wedge (A_n \vee D)$. Distribution gives us $C \vdash D$, which we already know not to be the case, since $\Sigma_n \not\vdash \Delta_n$ by hypothesis. So in this case too, $\Sigma_{n+1} \not\vdash \Delta_{n+1}$.

The final case is similar, except that we know that A_n has the form $\exists vB$, and we have $C \wedge A_n \vdash D$ and $C \vdash A_n \vee B(c) \vee D$. Conjoining C we get $C \vdash C \wedge (A_n \vee B(c) \vee D)$ for each c (and in particular, a c absent from C, A_n and D) and hence $C \vdash \forall v(C \wedge (A_n \vee B \vee D)) \vdash C \wedge (A_n \vee \forall vB \vee D) = C \wedge (A_n \vee D)$, and we proceed as before to get $C \vdash D$, which contradicts the assumption, giving then $\Sigma_{n+1} \not\vdash \Delta_{n+1}$.

The pair $\langle \Sigma', \Delta' \rangle$ is quantifier suited by construction too. If $\Sigma' \vdash \Delta' \cup \{A(c)\}$ for each constant c then we must have $A(c) \in \Sigma'$ for each c , and hence $\forall v A \in \Sigma'$, giving $\Sigma' \vdash \Delta' \cup \{\forall v A\}$. The existential quantifier case is perfectly dual. \triangleleft

Now once we have a domain, we want to *keep* it. This is where the rather weak definition of quantifier-suitedness comes in.

LEMMA 3 (PAIR EXTENSION LEMMA 2) *If $\langle \Sigma, \Delta \rangle$ is a quantifier-suited pair, in which one of Σ and Δ is finite, then there is a full quantifier-suited pair $\langle \Sigma', \Delta' \rangle$ extending $\langle \Sigma, \Delta \rangle$, in a the same language.*

To prove this lemma we need the following simple lemma.

LEMMA 4 (FINITE ADDITION LEMMA) *If $\langle \Sigma, \Delta \rangle$ is a quantifier-suited pair and X and Y are finite sets of formulas in the same language, and one of Σ and Δ is finite, then $\langle \Sigma \cup X, \Delta \cup Y \rangle$ is also quantifier-suited.*

This is the result for which we need subtraction and intuitionistic implication.

PROOF Here is the case where Σ is finite. By abuse of notation, I will let “ X ” and “ Y ” stand for the *formulae* $\bigwedge X$ and $\bigvee Y$. Let B be the formula $\bigwedge \Sigma$.

Now, if $\Sigma \cup X \cup \{A(c)\} \vdash \Delta \cup Y$ for each name c , we have $B \wedge X \wedge A(c) \vdash \Delta \cup Y$, and equivalently, $(B \wedge X \wedge A(c)) - Y \vdash \Delta$. So, by the quantifier-suitedness of $\langle \Sigma, \Delta \rangle$ we have $(\exists v)((B \wedge X \wedge A) - Y) \vdash \Delta$. As a result, $(B \wedge X \wedge \exists v A) - Y \vdash \Delta$, and hence $\Sigma \cup X \cup \{\exists v A\} \vdash \Delta \cup Y$ as desired.

If $\Sigma \cup X \vdash \Delta \cup Y \cup \{A(c)\}$ for each name c , we have $(B \wedge X) - (Y \vee A(c)) \vdash \Delta$ for each c , and hence $\exists v((B \wedge X) - (Y \vee A)) \vdash \Delta$. But this gives $(B \wedge X) - (Y \vee \forall v A) \vdash \Delta$ and hence $\Sigma \cup X \vdash \Delta \cup Y \cup \{\forall v A\}$ as desired.

The case for finite Δ is perfectly dual, using implication instead of subtraction. \triangleleft

Given the finite addition lemma, we can prove the second pair extension lemma.

PROOF The process is similar to that used in the proof of the first pair extension lemma. Now, however, instead of adding a *new* witness for each existential quantifier, we show that an old one will do.

- If $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$ and A_n is not of the form $\exists v B$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n \cup \{A_n\}, \Delta_n \rangle$.
- If $\Sigma_n \cup \{A_n\} \not\vdash \Delta_n$ and A_n is of the form $\exists v B$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n \cup \{A_n, B(c)\}, \Delta_n \rangle$, for some constant c such that $\Sigma_n \cup \{A_n, B[v := c]\} \not\vdash \Delta_n$.
- If $\Sigma_n \cup \{A_n\} \vdash \Delta_n$ and A_n is not of the form $\forall v B$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n, \Delta_n \cup \{A_n\} \rangle$.
- If $\Sigma_n \cup \{A_n\} \vdash \Delta_n$ and A_n is of the form $\forall v B$, then $\langle \Sigma_{n+1}, \Delta_{n+1} \rangle = \langle \Sigma_n, \Delta_n \cup \{A_n, B(c)\} \rangle$, for some constant c such that $\Sigma_n \not\vdash \Delta_n \cup \{A_n, B(c)\}$.

We need show that at each stage n , if we use the second or fourth lines of the definition, an appropriate c can be found. For this, we appeal to the finite addition lemma: Each $\langle \Sigma_n, \Delta_n \rangle$ is quantifier-suited, as it is a finite extension to the quantifier-suited $\langle \Sigma, \Delta \rangle$, and one of Σ and Δ is finite.

Now, suppose we are in the second case. If there is no suitable c such that $\Sigma_n \cup \{A_n, B[v := c]\} \not\vdash \Delta_n$. It follows that for each c , $\Sigma_n \cup \{A_n, B(c)\} \vdash \Delta_n$.

Therefore since Σ_n is quantifier suited, $\Sigma_n \cup \{A_n\} \vdash \Delta_n$. However, this is a contradiction. The fourth case is dual. \triangleleft

Now we must construct theories to use in the canonical model. The strategy is simple. If $\Sigma \not\vdash \Delta$, we extend $\langle \Sigma, \Delta \rangle$ to a full quantifier-suited $\langle \Sigma', \Delta' \rangle$. Then we use the class of full quantifier-suited pairs in this language as our canonical frame. Since they are full and quantifier-suited we know that these points interpret the extensional part of the language adequately. We must ensure that we interpret the modal operators well too. We define the accessibility relations in the obvious way.

- ΣR , if and only if for each A , if $\Box A \in \Sigma$ then $A \in \Sigma$.
- ΣS , if and only if for each A if $A \in \Sigma$, then $\Diamond A \in \Sigma$.
- ΣC , if and only if whenever $\sim A \in \Sigma$ $A \notin \Sigma$.
- $\Sigma D\Delta$ if and only if $A \notin \Sigma$, then $\neg A \in \Sigma$.

These conditions ensure that the accessibility relations satisfy the required interactions with the *ordering* on the canonical frame. This order is the *subsethood* relation on the theories.

These definitions of accessibility relations immediately ensure *half* of the evaluation conditions for modal operators. If $\Box A \in \Sigma$ then for each Σ , where ΣR , $A \in \Sigma$. To complete the picture we wish to show that if $\Box A \notin \Sigma$ then there is some Σ , where ΣR , and $A \notin \Sigma$. This is slightly more difficult than in the propositional case, for we need to not only construct the theory, but also ensure that it is quantifier-suited in the same language. To do this we use the following lemma. This is the point at which the Barcan Laws are used.

LEMMA 5 *If $\langle \Sigma, \Delta \rangle$ is full and quantifier suited, then*

- *If $\Box B \notin \Sigma$ then $\langle \Box^{-1}\Sigma, \{B\} \rangle$ is a quantifier-suited pair.*
- *If $\neg B \notin \Sigma$ then $\langle \{B\}, \neg^{-1}\Sigma \rangle$ is a quantifier-suited pair.*
- *If $\Diamond B \in \Sigma$, then $\langle \{B\}, \Diamond^{-1}\Delta \rangle$ is a quantifier-suited pair.*
- *If $\neg B \in \Sigma$, then $\langle \neg^{-1}\Delta, \{B\} \rangle$ is a quantifier-suited pair.*
- *If $A \supset B \notin \Sigma$, then $\langle \Sigma \cup \{A\}, \{B\} \rangle$ is a quantifier-suited pair.*
- *If $A - B \in \Sigma$, then $\langle \{A\}, \{B\} \cup \Delta \rangle$ is a quantifier-suited pair.*

PROOF Let's look at three cases: for \Box , \neg and subtraction. First, the \Box case: $\langle \Box^{-1}\Sigma, \{B\} \rangle$ is a pair iff $\bigwedge_i A_i \not\vdash B$ for any finite set of A_i s in $\Box^{-1}\Sigma$. But this is simple: if $\bigwedge_i A_i \vdash B$, then $\Box \bigwedge_i A_i \vdash \Box B$, and hence $\bigwedge_i \Box A_i \vdash \Box B$, giving $\Box B \in \Sigma$ (since each $\Box A_i \in \Sigma$) contrary to our assumption.

$\langle \Box^{-1}\Sigma, \{B\} \rangle$ is quantifier suited if $\langle \Box^{-1}\Sigma, \emptyset \rangle$ is, by the finite addition lemma. And for this we first need to show that if $\Box^{-1}\Sigma \vdash A(c)$ for each c then $\Box^{-1}\Sigma \vdash \forall v A$. So, if $\bigwedge_i A_i \vdash A(c)$ for each c , it follows that $\Box \bigwedge_i A_i \vdash \Box A(c)$ for each v , and hence, $\bigwedge_i \Box A_i \vdash \Box A(c)$, giving $\Sigma \vdash A(c)$ for each c . Since $\langle \Sigma, \Delta \rangle$ is quantifier suited, it follows that $\forall v \Box A \in \Sigma$. The Barcan formula then gives $\Box \forall v A \in \Sigma$. As a result, $\Box^{-1}\Sigma \vdash \forall v A$ as desired.

For the second half, it is sufficient to show that if $\Box^{-1}\Sigma \cup \{A(c)\} \vdash \perp$ for each c then $\Box^{-1}\Sigma \vdash A(c) \supset \perp$ for each c , so the previous result applies, giving us $\Box^{-1}\Sigma \vdash \forall v(A \supset \perp)$ and hence $\Box^{-1}\Sigma \vdash \exists vA \supset \perp$ giving $\Box^{-1}\Sigma \cup \{\exists vA\} \vdash \perp$ as desired.

Now for the \frown case. First we show that $\langle \frown^{-1}\Delta, \{B\} \rangle$ is a pair. If $\bigwedge_i A_i \vdash B$ where $A_i \in \frown^{-1}\Delta$, then by the O conditions, $\frown B \vdash \frown \bigwedge_i A_i$ but $\frown \bigwedge_i A_i \vdash \bigvee_i \frown A_i$. But $\frown B \in \Sigma$, and this gives $\Sigma \vdash \Delta$ contrary to our assumption. So $\langle \frown^{-1}\Delta, \{B\} \rangle$ is a pair.

To show that it is quantifier suited, it suffices to show that $\langle \frown^{-1}\Delta, \emptyset \rangle$ is quantifier suited. For this, suppose $\frown^{-1}\Delta \vdash C(c)$ for each c . We then have $\frown C(c) \vdash \frown \bigwedge_i A_i$ and hence $\frown C(c) \vdash \bigvee_i \frown A_i$ (for a choice of the A_i from $\frown^{-1}\Delta$). So, $\frown C(c) \vdash \Delta$ and by the quantifier suitedness of the original pair, $\exists v \frown C \vdash \Delta$. By the Barcan Law for \frown we get $\frown \forall v C \vdash \Delta$, and hence $\frown^{-1}\Delta \vdash \forall v C$ as desired.

For the second half, it is sufficient to show that if $\frown^{-1}\Delta \cup \{A(c)\} \vdash \perp$ for each c then $\frown^{-1}\Delta \vdash A(c) \supset \perp$ for each c and as before, the previous result applies, giving us $\frown^{-1}\Delta \vdash \forall v(A \supset \perp)$ and hence $\frown^{-1}\Delta \vdash \exists vA \supset \perp$ giving $\frown^{-1}\Delta \cup \{\exists vA\} \vdash \perp$ as desired.

Finally, for subtraction, we wish to show that $\langle \{A\}, \{B\} \cup \Delta \rangle$ is a pair if $A - B \notin \Sigma$. This is immediate, for if $A \vdash \{B\} \cup \Delta$, then $A - B \vdash \Delta$. To show that it is quantifier suited, we wish to show that $\langle \emptyset, \{B\} \cup \Delta \rangle$ is quantifier suited. For this, if $C(c) \vdash \{B\} \cup \Delta$ for each c , we have $C(c) - B \vdash \Delta$ for each c and by the quantifier suitedness of the original pair, $\exists v(C - B) \vdash \Delta$ and hence $\exists vC - B \vdash \Delta$ (the v can be chosen to be absent from B) giving $\exists vC \vdash \Delta \cup \{B\}$. Furthermore, if $\top \vdash \{C(c), B\} \cup \Delta$ for each c , we have $\top - C(c) \vdash \{B\} \cup \Delta$ for each c , giving us $\exists v(\top - C) \vdash \{B\} \cup \Delta$ and hence $\top \vdash \{\forall vC, B\} \cup \Delta$ as desired. \triangleleft

These lemmas jointly give us our completeness proof.

THEOREM 6 (COMPLETENESS) *If $\Sigma \not\vdash \Delta$ then there is some model M in which $\Sigma \not\vdash \Delta$.*

PROOF Extend $\langle \Sigma, \Delta \rangle$ to a full quantifier suited pair $\langle \Sigma', \Delta' \rangle$ by the first pair extension lemma. Construct a frame on the set of all full quantifier suited pairs on the language of $\langle \Sigma', \Delta' \rangle$. The domain of this frame is the set of constants in the language. The extension of a predicate F at the a point $\langle \cdot, \Pi \rangle$ is the set of n -tuples $\langle c_1, \dots, c_n \rangle$ such that $Fc_1 \cdots c_n \in \cdot$. The lemmas proved above ensure that all of the connective evaluation conditions hold in the canonical frame, such that $\langle \cdot, \Pi \rangle \vdash A$ if and only if $A \in \cdot$. \triangleleft

5 Miscellany

WHAT OF THE NEW CONNECTIVES? Much more should be said about the new connectives \supset and $-$. Can the finite addition lemma be proved without them? This would certainly be an advantage. Are these connectives *meaningful* in any logics for which we want constant domain frames?

RESTRICTED QUANTIFICATION: We can let $E!$ be an existence predicate, and define $(\forall_{E!x})A$ to be $(\forall x)(E!x \supset A)$, and $(\exists_{E!x})A$ to be $(\exists x)(E!x \wedge A)$. This mimics increasing domain logics in a familiar fashion.

More oddly, we can define $N!$ as a *nonexistence* predicate, to define $(\forall_{N!x})A$ as $(\forall x)(N!x \vee A)$, and $(\exists_{N!x})$ to be $(\exists x)(A - N!x)$. This mimics *decreasing* domain logics. Is there any use for this dualisation of the intuitionistic case?

WHAT ABOUT LOGICS WITHOUT DISTRIBUTION? None of this works in the absence of distribution of conjunction over disjunction. For an analagous semantics to the frames presented here, you should examine the linear logic literature on *phase spaces* [7, 11].

DUNN CONDITIONS: An obvious pair of conditions to want \Box and \Diamond to satisfy is what I have called the *Dunn Conditions* [4]:

$$\Box(A \vee B) \vdash \Box A \vee \Diamond B \quad \Diamond A \wedge \Box B \vdash \Diamond(A \wedge B)$$

(They should look reminiscent of the quantifier distribution laws.) These are valid if \Box and \Diamond use the same accessibility relation. To prove the addition sound and complete for the obvious class of frames we need a stronger version of the finite addition lemma. For given $\Box A \notin \Sigma$ (with a quantifier suited pair $\langle \Sigma, \Delta \rangle$) we need $\langle \Box^{-1}\Sigma, \{A\} \cup \Diamond^{-1}\Delta \rangle$ to be quantifier suited. At the new accessible point, everything necessary at Σ is true, and everything not possible at Σ is not true. The current proof of the finite addition lemma requires that one of the parts of the pair be finite. This does not hold here. To prove completeness we need a stronger version of the finite addition lemma or a completely different proof.

IMPLICATION AND FUSION: Implication and fusion seem to cause more difficulties too. For example, with implication, if $A \rightarrow B \notin \Sigma$ where $\langle \Sigma, \Delta \rangle$ is quantifier-suited, we need to construct *two* pairs, one at which A holds (that is simple: show that $\langle \{A\}, \{C : C \rightarrow B \in \Sigma\} \rangle$ is quantifier-suited and then extend that to a full quantifier suited pair $\langle \cdot, \Pi \rangle$) and another at which B fails. This one is more difficult — we wish to show that $\langle \Sigma, \{B\} \rangle$ is quantifier-suited, where $\Sigma, = \{C \circ D : C \in \Sigma, D \in \cdot\}$. How do you do this?

IDENTITY: There is no doubt that there are choices to be made in the semantics of identity. If the extension of the identity predicate is reflexive at all points, then we have $A \vdash a = a$ for any proposition A . This will be anathema to relevant logicians. To give purchase to a relevant account of identity, we must have points at which identity fails to be reflexive. The validity of $a = b \vdash b = a$ ensures that identity is *symmetric* at all points. If $a = b \wedge b = c \vdash a = c$, then identity must be *transitive* at all points.⁵

References

- [1] J. MICHAEL DUNN. “Gaggle Theory: An Abstraction of Galois Connections and Residuation with Applications to Negation and Various Logical Operations”. In *Logics in AI, Proceedings European Workshop JELIA 1990*, volume 478 of *Lecture Notes in Computer Science*. Springer Verlag, 1991.
- [2] J. MICHAEL DUNN. “Gaggle Theory Applied to Modal, Intuitionistic and Relevance Logics”. In I. MAX AND W. STELZNER, editors, *Logik und Mathematik: Frege-Kolloquium Jena*. de Gruyter, 1993.

⁵Thanks to an audience at the 1998 Australasian Association for Logic Conference, for helpful discussion on an earlier version of this paper.

- [3] J. MICHAEL DUNN. “Star and Perp: Two Treatments of Negation”. In JAMES E. TOMBERLIN, editor, *Philosophical Perspectives*, volume 7, pages 331–357. Ridgeview Publishing Company, Atascadero, California, 1994.
- [4] J. MICHAEL DUNN. “Positive Modal Logic”. *Studia Logica*, 55:301–317, 1995.
- [5] KIT FINE. “Semantics for Quantified Relevance Logic”. *Journal of Philosophical Logic*, 17:27–59, 1988.
- [6] KIT FINE. “Incompleteness for Quantified Relevance Logics”. In JEAN NORMAN AND RICHARD SYLVAN, editors, *Directions in Relevant Logic*, pages 205–225. Kluwer Academic Publishers, Dordrecht, 1989.
- [7] JEAN-YVES GIRARD. “Linear Logic”. *Theoretical Computer Science*, 50:1–101, 1987.
- [8] G. HUGHES AND M. CRESSWELL. *A New Introduction to Modal Logic*. Routledge, London, 1996.
- [9] BERNARD LINSKY AND EDWARD N. ZALTA. “In Defense of the Simplest Quantified Modal Logic”. *Philosophical Perspectives*, 8:431–458, 1994.
- [10] GREG RESTALL. “Extending Intuitionistic Logic with Subtraction”. In Progress. Draft available from <http://www.mq.edu.au/~phildept/staff/grestall/>.
- [11] GREG RESTALL. *Consequences: An Introduction to Substructural Logics*. to appear, 1998.
- [12] RICHARD ROUTLEY. “Problems and Solutions in the Semantics of Quantified Relevant Logics — I”. In A. I. ARRUDA, R. CHUAQUI, AND N. C. A. DA COSTA, editors, *Proceedings of the Fourth Latin American Symposium on Mathematical Logic*, pages 305–340. North-Holland, 1980.
- [13] RICHARD ROUTLEY AND ROBERT K. MEYER. “Semantics of Entailment”. In HUGUES LEBLANC, editor, *Truth Syntax and Modality*, pages 194–243. North-Holland, Amsterdam, 1973. Proceedings of the Temple University Conference on Alternative Semantics.